arXiv:0904.3434v1 [math-ph] 22 Apr 2009

Drinfeld-Sokolov hierarchies of type A and fourth order Painlevé systems

Kenta Fuji and Takao Suzuki
Department of Mathematics, Kobe University
Rokko, Kobe 657-8501, Japan
E-mail: suzukit@math.kobe-u.ac.jp

Abstract

We study the Drinfeld-Sokolov hierarchies of type $A_n^{(1)}$ associated with the regular conjugacy classes of $W(A_n)$. A class of fourth order Painlevé systems is derived from them by similarity reductions.

1 Introduction

Three types of fourth order Painlevé type ordinary differential equations have been studied [FS, NY1, S]. They are extensions of the Painlevé equations $P_{\mathrm{II}}, \dots, P_{\mathrm{VI}}$ and expressed as Hamiltonian systems

$$\mathcal{H}^{X_n^{(1)}}: \quad \frac{dq_i}{dt} = \frac{\partial H^{X_n^{(1)}}}{\partial p_i}, \quad \frac{dp_i}{dt} = -\frac{\partial H^{X_n^{(1)}}}{\partial q_i} \quad (i = 1, 2),$$

with the Coupled Hamiltonians

$$\begin{aligned}
H^{A_4^{(1)}} &= H_{\mathrm{IV}}(q_1, p_1; \alpha_2, \alpha_1) + H_{\mathrm{IV}}(q_2, p_2; \alpha_4, \alpha_1 + \alpha_3) + 2q_1p_1p_2, \\
tH^{A_5^{(1)}} &= H_{\mathrm{V}}(q_1, p_1; \alpha_2, \alpha_1, \alpha_1 + \alpha_3) \\
&\quad + H_{\mathrm{V}}(q_2, p_2; \alpha_4, \alpha_1 + \alpha_3, \alpha_1 + \alpha_3) + 2q_1p_1(q_2 - 1)p_2, \\
t(t-1)H^{D_6^{(1)}} &= H_{\mathrm{VI}}(q_1, p_1; \alpha_0, \alpha_3 + \alpha_5, \alpha_3 + \alpha_6, \alpha_2(\alpha_1 + \alpha_2)) \\
&\quad + H_{\mathrm{VI}}(q_2, p_2; \alpha_0 + \alpha_3, \alpha_5, \alpha_6, \alpha_4(\alpha_1 + 2\alpha_2 + \alpha_3 + \alpha_4)) \\
&\quad + 2(q_1 - t)p_1q_2\{(q_2 - 1)p_2 + \alpha_4\},
\end{aligned}$$

Lie algebra	Partition	Painlevé system
$A_1^{(1)}$	(2)	P_{II}
	$(1,1)$	P_{IV}
$A_2^{(1)}$	(3)	P_{IV}
	$(2,1)$	P_{V}
	$(1,1,1)$	P_{VI}
$A_3^{(1)}$	(4)	P_{V}
$A_4^{(1)}$	(5)	$\mathcal{H}^{A_4^{(1)}}$
$A_5^{(1)}$	(6)	$\mathcal{H}^{A_5^{(1)}}$

Table 1: Relation between $A_n^{(1)}$-hierarchies and Painlevé systems

where

$$\begin{aligned} H_{\mathrm{IV}}(q,p;a,b) &= qp(p-q-t) - aq - bp, \\ H_{\mathrm{V}}(q,p;a,b,c) &= q(q-1)p(p+t) + atq + bp - cqp, \\ H_{\mathrm{VI}}(q,p;a,b,c,d) &= q(q-1)(q-t)p^2 - \{(a-1)q(q-1) \\ &\quad + bq(q-t) + c(q-1)(q-t)\}p + dq. \end{aligned}$$

But complete classification of fourth order Painlevé systems is not achieved, so that the existence of unknown ones is expected. In this article, we derive a class of fourth order Painlevé systems from the Drinfeld-Sokolov hierarchies of type $A_n^{(1)}$ by similarity reductions.

The Drinfeld-Sokolov hierarchies are extensions of the KdV (or mKdV) hierarchy for the affine Lie algebras [DS]. For type $A_n^{(1)}$, they imply several Painlevé systems by similarity reductions [AS, KIK, KK1, KK2, NY1]; *see Table 1.* Such fact clarifies the origines of several properties of the Painlevé systems, Lax pairs, affine Weyl group symmetries and particular solutions in terms of the Schur polynomials.

The Drinfeld-Sokolov hierarchies are characterized by the Heisenberg subalgebras, that is maximal nilpotent subalgebras, of the affine Lie algebras. And the isomorphism classes of the Heisenberg subalgebras are in one-to-one correspondence with the conjugacy classes of the finite Weyl group [KP]. In this article, we choose the *regular* conjugacy classes of $W(A_n)$ and consider their associated hierarchies, called *type I hierarchies* [GHM]. In the notation of [DF], the regular conjugacy classes of $W(A_n)$ correspond to the partitions $(p,\dots,p)$ and $(p,\dots,p,1)$. For the derivation of fourth order Painlevé sys-

Lie algebra	Partition	Painlevé system
$A_3^{(1)}$	$(2,2)$	P_{VI}
	$(3,1)$	$\mathcal{H}^{A_4^{(1)}}$
$A_4^{(1)}$	$(4,1)$	$\mathcal{H}^{A_5^{(1)}}$
	$(2,2,1)$	system (1.1) with (1.2)
$A_5^{(1)}$	$(3,3)$	system (1.1) with (1.2)

Table 2: List of Painlevé systems obtained in this article

tems, we investigate the partitions $(2,2)$, $(3,1)$, $(4,1)$, $(2,2,1)$ and $(3,3)$; *see Table 2.*

One of impotant results in this article is the derivation of a new Painlevé system. It is expressed as a Hamiltonian system

$$\frac{dq_i}{dt} = \frac{\partial H_c}{\partial p_i}, \quad \frac{dp_i}{dt} = -\frac{\partial H_c}{\partial q_i} \quad (i = 1, 2), \tag{1.1}$$

with a Coupled Hamiltonian

$$\begin{aligned} t(t-1)H_c = {} & H_{\mathrm{VI}}(q_1, p_1; \alpha_2, \alpha_0 + \alpha_4, \alpha_3 + \alpha_5 - \eta, \eta\alpha_1) \\ & + H_{\mathrm{VI}}(q_2, p_2; \alpha_0 + \alpha_2, \alpha_4, \alpha_1 + \alpha_3 - \eta, \eta\alpha_5) \\ & + (q_1 - t)(q_2 - 1)\left\{(q_1p_1 + \alpha_1)p_2 + p_1(p_2q_2 + \alpha_5)\right\}. \end{aligned} \tag{1.2}$$

This system admits affine Weyl group symmetry of type $A_5^{(1)}$; see Appendix B. On the other hand, the system $\mathcal{H}^{D_6^{(1)}}$ admits one of type $D_6^{(1)}$. The relation between those two coupled Painlevé VI systems is not clarified.

Remark 1.1. *For the partition* $(1, \ldots, 1)$ *of* $n + 2$*, we have the Garnier system in* n*-variables* [KK2]. *Also for each partition* $(5, 1)$ *and* $(2, 2, 2)$*, a system of sixth order is derived*; *we do not give the explicit formula here. Thus we conjecture that any more fourth order Painlevé system do not arise from the type I hierarchy.*

This article is organized as follows. In Section 2, we recall the affine Lie algebra of type $A_n^{(1)}$ and realize it in a flamework of a central extension of the loop algebra $\mathfrak{sl}_{n+1}[z, z^{-1}]$. In Section 3, the Heisenberg subalgebra of $\widehat{\mathfrak{sl}}_{n+1}$ corresponding to the partition $\mathbf{n}$ is introduced. In Section 4, we formulate the Drinfeld-Sokolov hierarchies and their similarity reductions. In Section 5 and 6, the Painlevé systems are derived from the Drinfeld-Sokolov hierarchies. In

Appendix A, we give explicit descriptions of Lax pairs by means of a bases of $\widehat{\mathfrak{sl}}_{n+1}$. In Appendix B, we discuss a group of symmetries for the system (1.1) with (1.2).

2 Affine Lie algebra

In this section, we recall the affine Lie algebra of type $A_n^{(1)}$ and realize it in a flamework of a central extension of the loop algebra $\mathfrak{sl}_{n+1}[z, z^{-1}]$.

In the notation of [Kac], the affine Lie algebra $\mathfrak{g} = \mathfrak{g}(A_n^{(1)})$ is generated by the Chevalley generators $e_i, f_i, \alpha_i^\vee$ $(i = 0, \ldots, n)$ and the scaling element d with the fundamental relations

$$\begin{aligned}
&(\mathrm{ad}e_i)^{1-a_{i,j}}(e_j) = 0, \quad (\mathrm{ad}f_i)^{1-a_{i,j}}(f_j) = 0 \quad (i \neq j),\\
&[\alpha_i^\vee, \alpha_j^\vee] = 0, \quad [\alpha_i^\vee, e_j] = a_{i,j}e_j, \quad [\alpha_i^\vee, f_j] = -a_{i,j}f_j, \quad [e_i, f_j] = \delta_{i,j}\alpha_i^\vee,\\
&[d, \alpha_i^\vee] = 0, \quad [d, e_i] = \delta_{i,0}e_0, \quad [d, f_i] = -\delta_{i,0}f_0,
\end{aligned}$$

for $i, j = 0, \ldots, n$. The generalized Cartan matrix $A = [a_{i,j}]_{i,j=0}^n$ for $\mathfrak{g}$ is defined by

$$\begin{aligned}
&a_{i,i} = 2 && (i = 0, \ldots, n),\\
&a_{i,i+1} = a_{n,0} = a_{i+1,i} = a_{0,n} = -1 && (i = 0, \ldots, n-1),\\
&a_{i,j} = 0 && (\text{otherwise}).
\end{aligned}$$

We denote the Cartan subalgebra of $\mathfrak{g}$ by

$$\mathfrak{h} = \mathbb{C}\alpha_0^\vee \oplus \mathbb{C}\alpha_1^\vee \oplus \cdots \oplus \mathbb{C}\alpha_n^\vee \oplus \mathbb{C}d = \mathfrak{h}' \oplus \mathbb{C}d.$$

The normalized invariant form $(\cdot|\cdot) : \mathfrak{g} \times \mathfrak{g} \to \mathbb{C}$ is determined by the conditions

$$\begin{aligned}
&(\alpha_i^\vee|\alpha_j^\vee) = a_{i,j}, && (e_i|f_j) = \delta_{i,j}, && (\alpha_i^\vee|e_j) = (\alpha_i^\vee|f_j) = 0,\\
&(d|d) = 0, && (d|\alpha_j^\vee) = \delta_{0,j}, && (d|e_j) = (d|f_j) = 0,
\end{aligned}$$

for $i, j = 0, \ldots, n$.

Let $\mathfrak{n}_+$ and $\mathfrak{n}_-$ be the subalgebras of $\mathfrak{g}$ generated by e_i and f_i $(i = 0, \ldots, n)$ respectively. Then the Borel subalgebra $\mathfrak{b}_+$ of $\mathfrak{g}$ is defined by $\mathfrak{b}_+ = \mathfrak{h} \oplus \mathfrak{n}_+$. Note that we have the triangular decomposition

$$\mathfrak{g} = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}_+ = \mathfrak{n}_- \oplus \mathfrak{b}_+.$$

The corresponding infinite demensional groups are defined by

$$N_\pm = \exp(\mathfrak{n}_\pm^*), \quad H = \exp(\mathfrak{h}'), \quad B_+ = HN_+,$$

where $\mathfrak{n}_\pm^*$ are completions of $\mathfrak{n}_\pm$ respectively.

Let $\mathbf{s} = (s_0, \ldots, s_n)$ be a vector of non-negative integers. We consider a gradation $\mathfrak{g} = \bigoplus_{k\in\mathbb{Z}} \mathfrak{g}_k(\mathbf{s})$ of type $\mathbf{s}$ by setting

$$\deg \mathfrak{h} = 0, \quad \deg e_i = s_i, \quad \deg f_i = -s_i \quad (i = 0, \ldots, n).$$

With an element $\vartheta(\mathbf{s}) \in \mathfrak{h}$ such that

$$(\vartheta(\mathbf{s})|\alpha_i^\vee) = s_i \quad (i = 0, \ldots, n),$$

this gradation is defined by

$$\mathfrak{g}_k(\mathbf{s}) = \left\{ x \in \mathfrak{g} \;\middle|\; [\vartheta(\mathbf{s}), x] = kx \right\} \quad (k \in \mathbb{Z}).$$

We denote by

$$\mathfrak{g}_{<k}(\mathbf{s}) = \bigoplus_{l<k} \mathfrak{g}_l(\mathbf{s}), \quad \mathfrak{g}_{\geq k}(\mathbf{s}) = \bigoplus_{l\geq k} \mathfrak{g}_l(\mathbf{s}).$$

Note that a gradation $\mathbf{s}_p = (1, \ldots, 1)$, called *the principal gradation*, implies

$$\mathfrak{g}_{<0}(\mathbf{s}_p) = \mathfrak{n}_-, \quad \mathfrak{g}_{\geq 0}(\mathbf{s}_p) = \mathfrak{b}_+.$$

The affine Lie algebra $\mathfrak{g}$ can be identified with

$$\widehat{\mathfrak{sl}}_{n+1} = \mathfrak{sl}_{n+1}[z, z^{-1}] \oplus \mathbb{C}z\frac{d}{dz} \oplus \mathbb{C}K,$$

where K is a canonical central element. In a flamework of $\widehat{\mathfrak{sl}}_{n+1}$, the Chevalley generators and the scaling element are given by

$$\begin{aligned} &e_i = E_{i,i+1}, \quad f_i = E_{i+1,i}, \quad \alpha_i^\vee = E_{i,i} - E_{i+1,i+1} \quad (i = 1, \ldots, n), \\ &e_0 = zE_{n+1,1}, \quad f_0 = z^{-1}E_{1,n+1}, \quad \alpha_0^\vee = E_{n+1,n+1} - E_{1,1} + K, \quad d = z\frac{d}{dz}, \end{aligned}$$

where $E_{i,j} = (\delta_{i,r}\delta_{j,s})_{r,s=1}^{n+1}$ are matrix units. The Lie bracket is defined by

$$[z^k X, z^l Y] = z^{k+l}(XY - YX) + k\delta_{k+l,0}\mathrm{tr}(XY)K,$$

where $X, Y \in \mathfrak{sl}_{n+1}$.

3 Heisenberg subalgebra

For type $A_n^{(1)}$, the isomorphism classes of the Heisenberg subalgebras are in one-to-one correspondence with the partitions of $n+1$. In this section, we introduce the Heisenberg subalgebra of $\widehat{\mathfrak{sl}}_{n+1}$ corresponding to the partition $\mathbf{n}$ following the manner in [KL].

Let $\mathbf{n} = (n_1, n_2, \ldots, n_r, n_{r+1}, \ldots, n_s)$ be a partition of $n+1$ with $n_1 \geq n_2 \geq \ldots \geq n_r > n_{r+1} = \ldots = n_s = 1$. Consider a partition of matrix corresponding to $\mathbf{n}$

$$\begin{bmatrix} B_{11} & B_{12} & \cdots & B_{1s} \\ B_{21} & B_{22} & \cdots & B_{2s} \\ \vdots & \vdots & \ddots & \vdots \\ B_{s1} & B_{s2} & \cdots & B_{ss} \end{bmatrix},$$

where each block B_{ij} is an $n_i \times n_j$-matrix. With this blockform, we define matricies $\Lambda'_i \in \widehat{\mathfrak{sl}}_{n+1}$ $(i = 1, \ldots, r)$ by

$$\Lambda'_i = \begin{bmatrix} O & \cdots & O \\ & & \\ \vdots & B_{ii} & \vdots \\ & & \\ O & \cdots & O \end{bmatrix}, \quad B_{ii} = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & & 0 \\ \vdots & \vdots & & \ddots & \\ 0 & 0 & 0 & & 1 \\ z & 0 & 0 & \cdots & 0 \end{bmatrix},$$

diagonal matricies $H'_j \in \widehat{\mathfrak{sl}}_{n+1}$ $(i = j, \ldots, s-1)$ by

$$H'_j = n_{j+1} z^{-1} (\Lambda'_j)^{n_j} - n_j z^{-1} (\Lambda'_{j+1})^{n_{j+1}},$$

and a diagonal matrix $\eta'_{\mathbf{n}} \in \widehat{\mathfrak{sl}}_{n+1}$ by

$$B_{ii} = \frac{1}{2n_i} \mathrm{diag}(n_i - 1, n_i - 3, \ldots, -n_i + 1) \quad (i = 1, \ldots, r).$$

Denoting the matrix $\eta'_{\mathbf{n}}$ by $\mathrm{diag}(\eta'_1, \eta'_2, \ldots, \eta'_{n+1})$, we consider a permutation

$$\sigma = \begin{pmatrix} \eta'_1 & \eta'_2 & \cdots & \eta'_{n+1} \\ \eta_1 & \eta_2 & \cdots & \eta_{n+1} \end{pmatrix},$$

such that $\eta_1 \geq \eta_2 \geq \ldots \geq \eta_{n+1}$. This permutation can be lifted to the transformation σ acting on the matricies Λ'_i and H'_j. We set

$$\Lambda_i = \sigma(\Lambda'_i) \quad (i = 1, \ldots, r), \quad H_j = \sigma(H'_j) \quad (j = 1, \ldots, s-1).$$

Then the Heisenberg subalgebra of $\widehat{\mathfrak{sl}}_{n+1}$ corresponding to the partition $\mathbf{n}$ is defined by

$$\mathfrak{s}_{\mathbf{n}} = \bigoplus_{i=1}^{r} \bigoplus_{k\in\mathbb{Z}\setminus n_i\mathbb{Z}} \mathbb{C}\Lambda_i^k \oplus \bigoplus_{j=1}^{s-1} \bigoplus_{k\in\mathbb{Z}\setminus\{0\}} \mathbb{C}z^k H_j \oplus \mathbb{C}K.$$

Let $N'_{\mathbf{n}}$ be the least common multiple of $n_1, \ldots, n_s$. Also let

$$N_{\mathbf{n}} = \begin{cases} N'_{\mathbf{n}} & \text{if } N'_{\mathbf{n}}\left(\dfrac{1}{n_i} + \dfrac{1}{n_j}\right) \in 2\mathbb{Z} \text{ for } \forall(i,j) \\ 2N'_{\mathbf{n}} & \text{otherwise} \end{cases}.$$

We consider a operator corresponding to $\mathbf{n}$

$$\vartheta_{\mathbf{n}} = N_{\mathbf{n}}\left(z\frac{d}{dz} + \mathrm{ad}\eta_{\mathbf{n}}\right),$$

where $\eta_{\mathbf{n}} = \sigma(\eta'_{\mathbf{n}})$. Then the operator $\vartheta_{\mathbf{n}}$ implies a gradation $\mathbf{s} = (s_0, \ldots, s_n)$ as follows:

$$\vartheta_{\mathbf{n}}(e_i) = s_i e_i \quad (i = 0, \ldots, n).$$

Note that the Heisenberg subalgebra $\mathfrak{s}_{\mathbf{n}}$ admits the gradation $\mathbf{s}$ defined by $\vartheta_{\mathbf{n}}$.

4 Drinfeld-Sokolov hierarchy

In this section, we formulate the Drinfeld-Sokolov hierarchy associated with the Heisenberg subalgebra $\mathfrak{s}_{\mathbf{n}}$. Its similarity reduction is also formulated.

Let Λ_i and H_j be the generators for $\mathfrak{s}_{\mathbf{n}}$ given in Section 3. Introducing time variables $t_{i,k}$ $(i = 1, \ldots, r; k \in \mathbb{N})$, we consider an N_-B_+-valued function $G = G(t_{1,1}, t_{1,2}, \ldots)$ defined by

$$G = \exp\left(\sum_{i=1}^{r}\sum_{k=1}^{\infty} t_{i,k}\Lambda_i^k\right) G(0).$$

Here we assume the $\mathbf{n}$-reduced condition

$$t_{i,l} = 0 \quad (i = 1, \ldots, r; l \in n_i\mathbb{N}).$$

Then we have a system of partial differential equations

$$\partial_{i,k}(G) = \Lambda_i^k G \quad (i = 1, \ldots, r; k \in \mathbb{N}), \tag{4.1}$$

where $\partial_{i,k} = \partial/\partial t_{i,k}$ Via the trianglar decomposition

$$G = W^{-1}Z, \quad W \in N_-, \quad Z \in B_+,$$

the system (4.1) implies *a Sato equation*

$$\partial_{i,k}(W) = B_{i,k}W - W\Lambda_i^k \quad (i = 1, \ldots, r; k \in \mathbb{N}), \tag{4.2}$$

where $B_{i,k}$ stands for the b_+-component of $W\Lambda_i^k W^{-1}$. The compatibility condition of (4.2) gives the Drinfeld-Sokolov hierarchy

$$[\partial_{i,k} - B_{i,k}, \partial_{j,l} - B_{j,l}] = 0 \quad (i, j = 1, \ldots, r; k, l \in \mathbb{N}). \tag{4.3}$$

Under the system (4.2), we consider an equation

$$(\vartheta_{\mathbf{n}} - \mathrm{ad}\rho)(W) = \sum_{i=1}^{r} \sum_{k=1}^{\infty} d_i k t_{i,k} \partial_{i,k}(W), \tag{4.4}$$

where $d_i = \deg \Lambda_i$ $(i = 1, \ldots, r)$ and $\rho = \sum_{j=1}^{s-1} \rho_j H_j$. Note that each ρ_j is independent of time vatiables $t_{i,k}$. The compatibility condition of (4.2) and (4.4) gives

$$[\vartheta_{\mathbf{n}} - M, \partial_{i,k} - B_{i,k}] = 0 \quad (i = 1, \ldots, r; k \in \mathbb{N}), \tag{4.5}$$

where

$$M = \rho + \sum_{i=1}^{r} \sum_{k=1}^{\infty} d_i k t_{i,k} B_{i,k}.$$

We call the systems (4.3) and (4.5) a similarity reduction of the Drinfeld-Sokolov hierarchy.

Remark 4.1. *The similarity reduction can be regarded as the compatibility condition of a Lax form*

$$\partial_{i,k}(\Psi) = B_{i,k}\Psi \quad (i = 1, \ldots, r; k \in \mathbb{N}), \quad \vartheta_{\mathbf{n}}(\Psi) = M\Psi.$$

Here an N_-B_+-valued function Ψ is given by

$$\Psi = W \exp\left(\sum_{i=1}^{r} \sum_{k=1}^{\infty} t_{i,k} \Lambda_i^k\right).$$

5 Derivation of Coupled P_{VI}

In this section, we derive the Painlevé system (1.1) with (1.2) from the Drinfeld-Sokolov hierarchies for $\mathfrak{s}_{(3,3)}$ and $\mathfrak{s}_{(2,2,1)}$ by similarity reductions.

5.1 For the partition $(3,3)$

At first, we define the Heisenberg subalgebra $\mathfrak{s}_{(3,3)}$ of $\mathfrak{g}(A_5^{(1)})$. Let

$$\Lambda_1 = e_{1,2} + e_{3,4} + e_{5,0}, \quad \Lambda_2 = e_{0,1} + e_{2,3} + e_{4,5}, \quad H_1 = \alpha_1^\vee + \alpha_3^\vee + \alpha_5^\vee,$$

where

$$e_{i_1,i_2,\dots,i_{n-1},i_n} = \mathrm{ad}e_{i_1}\mathrm{ad}e_{i_2}\dots\mathrm{ad}e_{i_{n-1}}(e_{i_n}).$$

Then we have

$$\mathfrak{s}_{(3,3)} = \bigoplus_{k\in\mathbb{Z}\setminus 3\mathbb{Z}} \mathbb{C}\Lambda_1^k \oplus \bigoplus_{k\in\mathbb{Z}\setminus 3\mathbb{Z}} \mathbb{C}\Lambda_2^k \oplus \bigoplus_{k\in\mathbb{Z}\setminus\{0\}} \mathbb{C}z^k H_1 \oplus \mathbb{C}K.$$

The grade operator for $\mathfrak{s}_{(3,3)}$ is given by

$$\vartheta_{(3,3)} = 3\left(z\frac{d}{dz} + \mathrm{ad}\eta_{(3,3)}\right),$$

where

$$\eta_{(3,3)} = \frac{1}{3}(\alpha_1^\vee + 2\alpha_2^\vee + 2\alpha_3^\vee + 2\alpha_4^\vee + \alpha_5^\vee).$$

It follows that $\mathfrak{s}_{(3,3)}$ admits the gradation of type $\mathbf{s} = (1,0,1,0,1,0)$, namely

$$\vartheta_{(3,3)}(e_i) = e_i \quad (i = 0,2,4), \quad \vartheta_{(3,3)}(e_j) = 0 \quad (j = 1,3,5).$$

Note that

$$\mathfrak{g}_{\geq 0}(1,0,1,0,1,0) = \mathbb{C}f_1 \oplus \mathbb{C}f_3 \oplus \mathbb{C}f_5 \oplus \mathfrak{b}_+.$$

We now assume $t_{2,1} = 1$ and $t_{1,k} = t_{2,k} = 0$ $(k \geq 2)$. Then the similarity reduction (4.3) and (4.5) for $\mathfrak{s}_{(3,3)}$ is expressed as

$$\left[\vartheta_{(3,3)} - M, \partial_{1,1} - B_{1,1}\right] = 0. \tag{5.1}$$

Here the $\mathfrak{b}_+$-valued functions M and $B_{1,1}$ are defined by

$$\begin{aligned} M &= \vartheta_{(3,3)}(W)W^{-1} + W(\rho_1 H_1 + t_{1,1}\Lambda_1 + \Lambda_2)W^{-1}, \\ B_{1,1} &= \partial_{1,1}(W)W^{-1} + W\Lambda_1 W^{-1}, \end{aligned} \tag{5.2}$$

where W is an N_--valued function; its explicit formula is given below. In the following, we derive the Painlevé system from the system (5.1) with (5.2).

We denote by

$$W = \exp(\omega_0)\exp(\omega_{-1})\exp(\omega_{<-1}),$$

where

$$\begin{aligned}\omega_0 &= -w_1 f_1 - w_3 f_3 - w_5 f_5,\\ \omega_{-1} &= -w_0 f_0 - w_2 f_2 - w_4 f_4 - w_{0,1} f_{0,1} - w_{1,2} f_{1,2} - w_{2,3} f_{2,3} - w_{3,4} f_{3,4}\\ &\quad - w_{4,5} f_{4,5} - w_{5,0} f_{5,0} - w_{1,2,3} f_{1,2,3} - w_{3,4,5} f_{3,4,5} - w_{5,0,1} f_{5,0,1},\end{aligned}$$

and $\omega_{<-1} \in \mathfrak{g}_{<-1}(1,0,1,0,1,0)$. Then the $\mathfrak{b}_+$-valued function M is described as

$$\begin{aligned}M &= \kappa_0 \alpha_0^\vee + \kappa_1 \alpha_1^\vee + \kappa_2 \alpha_2^\vee + \kappa_3 \alpha_3^\vee + \kappa_4 \alpha_4^\vee + \kappa_5 \alpha_5^\vee - (t_{1,1} w_5 - w_1) e_0 + \varphi_1 e_1\\ &\quad - (t_{1,1} w_1 - w_3) e_2 + \varphi_3 e_3 - (t_{1,1} w_3 - w_5) e_4 + \varphi_5 e_5 + t_{1,1} \Lambda_1 + \Lambda_2,\end{aligned}$$

with dependent variables

$$\varphi_1 = t_{1,1} w_2 - w_0, \quad \varphi_3 = t_{1,1} w_4 - w_2, \quad \varphi_5 = t_{1,1} w_0 - w_4,$$

and parameters

$$\begin{aligned}\kappa_0 &= -t_{1,1} w_{5,0} - w_{0,1}, & \kappa_1 &= t_{1,1}(w_1 w_2 - w_{1,2}) - (w_0 w_1 + w_{0,1}) + \rho_1,\\ \kappa_2 &= -t_{1,1} w_{1,2} - w_{2,3}, & \kappa_3 &= t_{1,1}(w_3 w_4 - w_{3,4}) - (w_2 w_3 + w_{2,3}) + \rho_1,\\ \kappa_4 &= -t_{1,1} w_{3,4} - w_{4,5}, & \kappa_5 &= t_{1,1}(w_0 w_5 - w_{5,0}) - (w_4 w_5 + w_{4,5}) + \rho_1.\end{aligned}$$

Note that

$$\partial_{1,1}(\kappa_i) = 0 \quad (i = 0, \ldots, 5).$$

We also remark that

$$w_1 \varphi_1 + w_3 \varphi_3 + w_5 \varphi_5 + \kappa_0 - \kappa_1 + \kappa_2 - \kappa_3 + \kappa_4 - \kappa_5 + 3\rho_1 = 0.$$

The $\mathfrak{b}_+$-valued function $B_{1,1}$ is described as

$$\begin{aligned}B_{1,1} &= u_0 K + (u_1 + w_1 x_1) \alpha_1^\vee + u_2 \alpha_2^\vee + (u_3 + w_3 x_3) \alpha_3^\vee + u_4 \alpha_4^\vee\\ &\quad + w_5 x_5 \alpha_5^\vee - w_5 e_0 + x_1 e_1 - w_1 e_2 + x_3 e_3 - w_3 e_4 + x_5 e_5 + \Lambda_1,\end{aligned}$$

where

$$\begin{aligned}u_1 &= \frac{-2w_1 \varphi_1 + w_3 \varphi_3 + w_5 \varphi_5 - 2\kappa_0 + 2\kappa_1 + \kappa_2 - \kappa_3 + \kappa_4 - \kappa_5}{3t_{1,1}},\\ u_2 &= -\frac{w_1 \varphi_1 + \kappa_0 - \kappa_1 + \rho_1}{t_{1,1}},\\ u_3 &= \frac{-w_1 \varphi_1 - w_3 \varphi_3 + 2w_5 \varphi_5 - \kappa_0 + \kappa_1 - \kappa_2 + \kappa_3 + 2\kappa_4 - 2\kappa_5}{3t_{1,1}},\\ u_4 &= \frac{w_5 \varphi_5 + \kappa_4 - \kappa_5 + \rho_1}{t_{1,1}}, \quad x_1 = \frac{t_{1,1}^2 \varphi_1 + t_{1,1} \varphi_5 + \varphi_3}{t_{1,1}^3 - 1},\\ x_3 &= \frac{t_{1,1}^2 \varphi_3 + t_{1,1} \varphi_1 + \varphi_5}{t_{1,1}^3 - 1}, \quad x_5 = \frac{t_{1,1}^2 \varphi_5 + t_{1,1} \varphi_3 + \varphi_1}{t_{1,1}^3 - 1}.\end{aligned}$$

Hence the system (5.1) with (5.2) can be expressed as a system of ordinary differential equations in terms of the variabes $\varphi_1, \varphi_5, w_1, w_3, w_5$; we do not give its explicit formula.

Let

$$q_1 = \frac{w_1}{t_{1,1}^2 w_3}, \quad p_1 = \frac{t_{1,1}^2 w_3 \varphi_1}{3}, \quad q_2 = \frac{w_5}{t_{1,1} w_3}, \quad p_2 = \frac{t_{1,1} w_3 \varphi_5}{3}, \quad t = \frac{1}{t_{1,1}^3}.$$

We also set

$$\begin{aligned}
\alpha_0 &= \frac{1}{3}(1 - 2\kappa_0 + \kappa_1 + \kappa_5), \quad &\alpha_1 &= \frac{1}{3}(\kappa_0 - 2\kappa_1 + \kappa_2), \\
\alpha_2 &= \frac{1}{3}(1 + \kappa_1 - 2\kappa_2 + \kappa_3), \quad &\alpha_3 &= \frac{1}{3}(\kappa_2 - 2\kappa_3 + \kappa_4), \\
\alpha_4 &= \frac{1}{3}(1 + \kappa_3 - 2\kappa_4 + \kappa_5), \quad &\alpha_5 &= \frac{1}{3}(\kappa_0 + \kappa_4 - 2\kappa_5),
\end{aligned}$$

and

$$\eta = \rho_1 + \frac{1}{2}(\alpha_1 + \alpha_3 + \alpha_5).$$

Then we have

Theorem 5.1. *The system* (5.1) *with* (5.2) *gives the Painlevé system* (1.1) *with* (1.2). *Furthermore, w_3 satisfies the completely integrable Pfaffian equation*

$$\begin{aligned}
t(t-1)\frac{d}{dt}\log w_3 = &-(q_1 - 1)(q_1 - t)p_1 - (q_2 - 1)(q_2 - t)p_2 \\
&- \alpha_1 q_1 - \alpha_5 q_2 + \frac{1}{3}(\alpha_1 + \alpha_2 - \alpha_3 - \alpha_4 + 2\eta)t \\
&- \frac{1}{3}(\alpha_1 + \alpha_2 + 2\alpha_3 - \alpha_4 - 4\eta).
\end{aligned}$$

5.2 For the partition $(2, 2, 1)$

The Heisenberg subalgebra $\mathfrak{s}_{(2,2,1)}$ of $\mathfrak{g}(A_4^{(1)})$ is defined by

$$\mathfrak{s}_{(2,2,1)} = \bigoplus_{k \in \mathbb{Z} \setminus 2\mathbb{Z}} \mathbb{C}\Lambda_1^k \oplus \bigoplus_{k \in \mathbb{Z} \setminus 2\mathbb{Z}} \mathbb{C}\Lambda_2^k \oplus \bigoplus_{k \in \mathbb{Z} \setminus \{0\}} \mathbb{C}z^k H_1 \oplus \bigoplus_{k \in \mathbb{Z} \setminus \{0\}} \mathbb{C}z^k H_2 \oplus \mathbb{C}K,$$

with

$$\begin{aligned}
\Lambda_1 &= e_{4,0} + e_{1,2,3}, \quad &\Lambda_2 &= e_{0,1} + e_{2,3,4}, \\
H_1 &= \alpha_1^\vee + \alpha_2^\vee - \alpha_3^\vee, \quad &H_2 &= -\alpha_2^\vee + \alpha_3^\vee + \alpha_4^\vee.
\end{aligned}$$

The subalgebra $\mathfrak{s}_{(2,2,1)}$ admits the gradation of type $\mathbf{s} = (2,0,1,1,0)$ with the grade operator

$$\vartheta_{(2,2,1)} = 4\left(z\frac{d}{dz} + \mathrm{ad}\eta_{(2,2,1)}\right), \quad \eta_{(2,2,1)} = \frac{1}{4}(\alpha_1^\vee + 2\alpha_2^\vee + 2\alpha_3^\vee + \alpha_4^\vee).$$

Note that

$$\mathfrak{g}_{\geq 0}(2,0,1,1,0) = \mathbb{C}f_1 \oplus \mathbb{C}f_4 \oplus \mathfrak{b}_+.$$

We now assume $t_{1,2} = 1$ and $t_{1,k} = t_{2,k} = 0$ $(k \geq 3)$. Then the similarity reduction (4.5) for $\mathfrak{s}_{(2,2,1)}$ is expressed as

$$\left[\vartheta_{(2,2,1)} - M, \partial_{1,1} - B_{1,1}\right] = 0, \tag{5.3}$$

with

$$\begin{aligned} M &= \vartheta_{(2,2,1)}(W)W^{-1} + W(\rho_1 H_1 + \rho_2 H_2 + 2t_{1,1}\Lambda_1 + 2\Lambda_2)W^{-1}, \\ B_{1,1} &= \partial_{1,1}(W)W^{-1} + W\Lambda_1 W^{-1}. \end{aligned} \tag{5.4}$$

Let

$$W = \exp(\omega_0)\exp(\omega_{-1})\exp(\omega_{-2})\exp(\omega_{<-2}),$$

where

$$\begin{aligned} \omega_0 &= -w_1 f_1 - w_4 f_4, \\ \omega_{-1} &= -w_2 f_2 - w_3 f_3 - w_{1,2} f_{1,2} - w_{3,4} f_{3,4}, \\ \omega_{-2} &= -w_0 f_0 - w_{0,1} f_{0,1} - w_{2,3} f_{2,3} - w_{4,0} f_{4,0} \\ &\quad - w_{1,2,3} f_{1,2,3} - w_{2,3,4} f_{2,3,4} - w_{4,0,1} f_{4,0,1} - w_{1,2,3,4} f_{1,2,3,4}, \end{aligned}$$

and $\omega_{<-2} \in \mathfrak{g}_{<-2}(2,0,1,1,0)$. Then the system (5.4) gives explicit formulas of $M, B_{1,1}$ as follows:

$$\begin{aligned} M &= \kappa_0\alpha_0^\vee + \kappa_1\alpha_1^\vee + \kappa_2\alpha_2^\vee + \kappa_3\alpha_3^\vee + \kappa_4\alpha_4^\vee + 2(w_1 - t_{1,1}w_4)e_0 \\ &\quad + \varphi_1 e_1 + (\varphi_2 - w_1\varphi_{1,2})e_2 + (\varphi_3 + w_4\varphi_{3,4})e_3 + \varphi_4 e_4 \\ &\quad + \varphi_{1,2}e_{1,2} + 2(t_{1,1}w_1 - w_4)e_{2,3} - \varphi_{3,4}e_{3,4} + 2t_{1,1}\Lambda_1 + 2\Lambda_2, \\ B_{1,1} &= u_0 K + (u_2 + w_1 x_1)\alpha_1^\vee + u_2\alpha_2^\vee + u_3\alpha_3^\vee + w_4 x_4\alpha_4^\vee - w_4 e_0 \\ &\quad + x_1 e_1 - w_1 x_{1,2} e_2 + \frac{\varphi_3}{2t_{1,1}} e_3 + x_4 e_4 + x_{1,2} e_{1,2} - w_1 e_{2,3} + \Lambda_1, \end{aligned}$$

where

$$\begin{aligned} &\varphi_1 = -2w_0 + t_{1,1}w_2 w_3 - 2t_{1,1}w_{2,3}, \quad \varphi_2 = -2w_{3,4}, \quad \varphi_3 = 2t_{1,1}w_{1,2}, \\ &\varphi_4 = 2t_{1,1}w_0 + w_2 w_3 + 2w_{2,3}, \quad \varphi_{1,2} = 2t_{1,1}w_3, \quad \varphi_{3,4} = -2w_2, \end{aligned}$$

and

$$
\begin{aligned}
u_2 &= -\frac{w_1\varphi_1+\kappa_0-\kappa_1+\rho_1}{2t_{1,1}}, \quad u_3 = \frac{w_4\varphi_4+\kappa_3-\kappa_4+\rho_1}{2t_{1,1}},\\
x_1 &= \frac{(t_{1,1}\varphi_1+\varphi_4)\varphi_3+(w_1\varphi_1+w_4\varphi_4+\kappa_0-\kappa_1+\kappa_3-\kappa_4+2\rho_1)\varphi_{3,4}}{2(t_{1,1}^2-1)\varphi_3},\\
x_4 &= \frac{(\varphi_1+t_{1,1}\varphi_4)\varphi_3+t_{1,1}(w_1\varphi_1+w_4\varphi_4+\kappa_0-\kappa_1+\kappa_3-\kappa_4+2\rho_1)\varphi_{3,4}}{2(t_{1,1}^2-1)\varphi_3},\\
x_{1,2} &= \frac{w_1\varphi_1+w_4\varphi_4+\kappa_0-\kappa_1+\kappa_3-\kappa_4+2\rho_1}{\varphi_3}.
\end{aligned}
$$

Note that $\kappa_0, \ldots, \kappa_4$ are constants. We also remark that

$$
\begin{aligned}
&\varphi_2\varphi_{3,4}+2(w_1\varphi_1+w_4\varphi_4+\kappa_0-\kappa_1+\kappa_2-\kappa_4+2\rho_2)=0,\\
&\varphi_3\varphi_{1,2}-2t_{1,1}(w_1\varphi_1+w_4\varphi_4+\kappa_0-\kappa_1+\kappa_3-\kappa_4+2\rho_1)=0.
\end{aligned}
$$

Hence the system (5.3) can be expressed as a system of ordinary differential equations in terms of the variables $\varphi_1, \varphi_3, \varphi_4, \varphi_{3,4}, w_1, w_4$.

Let

$$
\begin{aligned}
q_1 &= -\frac{t_{1,1}^2\varphi_{3,4}w_4}{\varphi_3}, \quad p_1 = -\frac{\varphi_3\varphi_4}{4t_{1,1}^2\varphi_{3,4}},\\
q_2 &= -\frac{t_{1,1}\varphi_{3,4}w_1}{\varphi_3}, \quad p_2 = -\frac{\varphi_3\varphi_1}{4t_{1,1}\varphi_{3,4}}, \quad t = t_{1,1}^2.
\end{aligned}
$$

We also set

$$
\begin{aligned}
\alpha_0 &= \frac{1}{4}(2-2\kappa_0+\kappa_1+\kappa_4), \quad \alpha_1 = \frac{1}{4}(\kappa_0+\kappa_3-2\kappa_4),\\
\alpha_2 &= \frac{1}{4}(1+\kappa_2-2\kappa_3+\kappa_4), \quad \alpha_3 = \frac{1}{4}(-\kappa_2+\kappa_3+2\rho_1-2\rho_2),\\
\alpha_4 &= \frac{1}{4}(1+\kappa_1-\kappa_2-2\rho_1+2\rho_2), \quad \alpha_5 = \frac{1}{4}(\kappa_0-2\kappa_1+\kappa_2),\\
\eta &= \frac{1}{4}(2\kappa_0-2\kappa_1+2\kappa_3-2\kappa_4+3\rho_1-\rho_2).
\end{aligned}
$$

Then we have

Theorem 5.2. *The system* (5.3) *with* (5.4) *gives the Painlevé system* (1.1) *with* (1.2). *Furthermore,* φ_3 *and* $\varphi_{3,4}$ *satisfy the completely integrable Pfaffian* *equations*

$$\begin{aligned}t(t-1)\frac{d}{dt}\log\varphi_3 &= -q_1(q_1-t)p_1-q_2(q_2-t)p_2-\alpha_1q_1-\alpha_5q_2\\&\quad+\frac{1}{4}(1+2\alpha_2-2\alpha_3-2\alpha_4-2\alpha_5+6\eta)t\\&\quad-\frac{1}{4}(1+2\alpha_2+2\alpha_3-2\alpha_4-2\alpha_5+2\eta),\\t(t-1)\frac{d}{dt}\log\varphi_{3,4} &= -(q_1-t)p_1-(q_2-t)p_2-\eta.\end{aligned}$$

6 Derivation of other systems

In this section, we discuss the derivation of the Painlevé systems for $\mathfrak{s}_{(2,2)}$, $\mathfrak{s}_{(3,1)}$ and $\mathfrak{s}_{(4,1)}$ by a similar manner as in Section 5.

6.1 For the partition $(2,2)$

The Heisenberg subalgebra $\mathfrak{s}_{(2,2)}$ of $\mathfrak{g}(A_3^{(1)})$ is defined by

$$\mathfrak{s}_{(2,2)}=\bigoplus_{k\in\mathbb{Z}\setminus2\mathbb{Z}}\mathbb{C}\Lambda_1^k\oplus\bigoplus_{k\in\mathbb{Z}\setminus2\mathbb{Z}}\mathbb{C}\Lambda_2^k\oplus\bigoplus_{k\in\mathbb{Z}\setminus\{0\}}\mathbb{C}z^kH_1\oplus\mathbb{C}K,$$

with

$$\Lambda_1=e_{1,2}+e_{3,0},\quad \Lambda_2=e_{0,1}+e_{2,3},\quad H_1=\alpha_1^\vee+\alpha_3^\vee.$$

The subalgebra $\mathfrak{s}_{(2,2)}$ admits the gradation of type $\mathbf{s}=(1,0,1,0)$ with the grade operator

$$\vartheta_{(2,2)}=2\left(z\frac{d}{dz}+\mathrm{ad}\eta_{(2,2)}\right),\quad \eta_{(2,2)}=\frac{1}{2}(\alpha_1^\vee+2\alpha_2^\vee+\alpha_3^\vee).$$

Note that

$$\mathfrak{g}_{\geq0}(1,0,1,0)=\mathbb{C}f_1\oplus\mathbb{C}f_3\oplus\mathfrak{b}_+.$$

We now assume $t_{1,2}=1$ and $t_{1,k}=t_{2,k}=0$ $(k\geq3)$. Then the similarity reduction (4.5) for $\mathfrak{s}_{(2,2)}$ is expressed as

$$\left[\vartheta_{(2,2)}-M,\partial_{1,1}-B_{1,1}\right]=0, \tag{6.1}$$

with

$$\begin{aligned}M&=\vartheta_{(2,2)}(W)W^{-1}+W(\rho_1H_1+t_{1,1}\Lambda_1+\Lambda_2)W^{-1},\\B_{1,1}&=\partial_{1,1}(W)W^{-1}+W\Lambda_1W^{-1}.\end{aligned} \tag{6.2}$$

Let

$$W = \exp(\omega_0)\exp(\omega_{-1})\exp(\omega_{<-1}),$$

where

$$\begin{aligned}\omega_0 &= -w_1 f_1 - w_3 f_3,\\ \omega_{-1} &= -w_0 f_0 - w_2 f_2 - w_{0,2} f_{0,2} - w_{1,2} f_{1,2}\\ &\quad - w_{2,3} f_{2,3} - w_{3,0} f_{3,0} - w_{1,2,3} f_{1,2,3} - w_{3,0,1} f_{3,0,1},\end{aligned}$$

and $\omega_{<-1} \in \mathfrak{g}_{<-1}(1,0,1,0)$. Then the system (6.2) gives explicit formulas of $M, B_{1,1}$ as follows:

$$\begin{aligned}M &= \kappa_0\alpha_0^\vee + \kappa_1\alpha_1^\vee + \kappa_2\alpha_2^\vee + \kappa_3\alpha_3^\vee + (w_1 - t_{1,1}w_3)e_0\\ &\quad + \varphi_1 e_1 + (w_3 - t_{1,1}w_1)e_2 + \varphi_3 e_3 + t_{1,1}\Lambda_1 + \Lambda_2,\\ B_{1,1} &= u_0 K + u_1\alpha_1^\vee + u_2\alpha_2^\vee + w_3 x_3\alpha_3^\vee + w_1 e_0 + x_1 e_1 + w_3 e_2 + x_3 e_3 + \Lambda_1,\end{aligned}$$

where

$$\varphi_1 = t_{1,1}w_2 - w_0, \quad \varphi_3 = t_{1,1}w_0 - w_2,$$

and

$$\begin{aligned}u_1 &= \frac{w_1}{t_{1,1}}x_3 - \frac{\kappa_0 - \kappa_1 + \rho_1}{t_{1,1}}, \quad u_2 = \frac{w_3\varphi_3 + \kappa_2 - \kappa_3 + \rho_1}{t_{1,1}},\\ x_1 &= \frac{(w_1 - t_{1,1}w_3)\varphi_3 - (\kappa_0 - \kappa_1 + \kappa_2 - \kappa_3 + 2\rho_1)t_{1,1}}{(t_{1,1}^2 - 1)w_1},\\ x_3 &= \frac{(t_{1,1}w_1 - w_3)\varphi_3 - (\kappa_0 - \kappa_1 + \kappa_2 - \kappa_3 + 2\rho_1)}{(t_{1,1}^2 - 1)w_1}.\end{aligned}$$

Note that $\kappa_0, \ldots, \kappa_3$ are constants. We also remark that

$$w_1\varphi_1 + w_3\varphi_3 + \kappa_0 - \kappa_1 + \kappa_2 - \kappa_3 + 2\rho_1 = 0.$$

Hence the system (6.1) can be expressed as a system of ordinary differential equations in terms of the variables φ_3, w_1, w_3.

Let

$$p = \frac{w_1\varphi_3}{2t_{1,1}}, \quad q = \frac{t_{1,1}w_3}{w_1}, \quad t = t_{1,1}^2.$$

We also set

$$\begin{aligned}&\alpha_0 = \frac{1}{2}(1 + \kappa_1 - 2\kappa_2 + \kappa_3), \quad \alpha_1 = \frac{1}{2}(-\kappa_1 + \kappa_3 + 2\rho_1),\\ &\alpha_2 = \kappa_0 + \kappa_2 - 2\kappa_3, \quad \alpha_3 = \frac{1}{2}(1 - 2\kappa_0 + \kappa_1 + \kappa_3),\\ &\alpha_4 = \frac{1}{2}(-\kappa_1 + \kappa_3 - 2\rho_1),\end{aligned}$$

and

$$a = \alpha_0, \quad b = \alpha_3, \quad c = \alpha_4, \quad d = \alpha_2(\alpha_1 + \alpha_2).$$

Then we have

Theorem 6.1. *The system* (6.1) *with* (6.2) *gives the sixth Painlevé equation. Furthermore, w_1 satisfies the completely integrable Pfaffian equation*

$$\begin{aligned} t(t-1)\frac{d}{dt}\log w_1 = &-(q-1)(q-t)p - \alpha_2 q \\ &+ \frac{1}{4}(1 + 2\alpha_1 - 2\alpha_3 - 4\alpha_4)t - \frac{1}{4}(1 - 2\alpha_1 - 4\alpha_2 - 2\alpha_3). \end{aligned}$$

6.2 For the partition $(3,1)$

The Heisenberg subalgebra $\mathfrak{s}_{(3,1)}$ of $\mathfrak{g}(A_3^{(1)})$ is defined by

$$\mathfrak{s}_{(3,1)} = \bigoplus_{k \in \mathbb{Z} \setminus 3\mathbb{Z}} \mathbb{C}\Lambda_1^k \oplus \bigoplus_{k \in \mathbb{Z} \setminus \{0\}} \mathbb{C}z^k H_1 \oplus \mathbb{C}K,$$

with

$$\Lambda_1 = e_0 + e_1 + e_{2,3}, \quad H_1 = \alpha_1^\vee + 2\alpha_2^\vee - \alpha_3^\vee.$$

The subalgebra $\mathfrak{s}_{(3,1)}$ admits the gradation of type $\mathbf{s} = (1,1,0,1)$ with the grade operator

$$\vartheta_{(3,1)} = 3z\left(\frac{d}{dz} + \mathrm{ad}\eta_{(3,1)}\right), \quad \eta_{(3,1)} = \frac{1}{3}(\alpha_1^\vee + \alpha_2^\vee + \alpha_3^\vee).$$

Note that

$$\mathfrak{g}_{\geq 0}(1,1,0,1) = \mathbb{C}f_2 \oplus \mathfrak{b}_+.$$

We now assume $t_{1,2} = 1$ and $t_{1,k} = 0$ ($k \geq 3$). Then the similarity reduction (4.5) for $\mathfrak{s}_{(3,1)}$ is expressed as

$$\left[\vartheta_{(3,1)} - M, \partial_{1,1} - B_{1,1}\right] = 0, \tag{6.3}$$

with

$$\begin{aligned} M &= \vartheta_{(3,1)}(W)W^{-1} + W(\rho_1 H_1 + t_{1,1}\Lambda_1 + 2\Lambda_1^2)W^{-1}, \\ B_{1,1} &= \partial_{1,1}(W)W^{-1} + W\Lambda_1 W^{-1}. \end{aligned} \tag{6.4}$$

Let

$$W = \exp(-w_2 f_2)\exp(\omega_{-1})\exp(\omega_{-2})\exp(\omega_{<-2}),$$

where

$$\begin{aligned}\omega_{-1} &= -w_0 f_0 - w_1 f_1 - w_3 f_3 - w_{1,2} f_{1,2} - w_{2,3} f_{2,3},\\ \omega_{-2} &= -w_{0,1} f_{0,1} - w_{3,0} f_{3,0} - w_{0,1,2} f_{0,1,2} - w_{1,2,3} f_{1,2,3} - w_{2,3,0} f_{2,3,0},\end{aligned}$$

and $\omega_{<-2} \in \mathfrak{g}_{<-2}(1,1,0,1)$. Then the system (6.4) gives explicit formulas of $M, B_{1,1}$ as follows:

$$\begin{aligned}M &= \kappa_0 \alpha_0^\vee + \kappa_1 \alpha_1^\vee + \kappa_2 \alpha_2^\vee + \kappa_3 \alpha_3^\vee + \varphi_0 e_0 + (\varphi_1 + w_2 \varphi_{1,2}) e_1\\ &\quad + \varphi_2 e_2 + (\varphi_3 - w_2 \varphi_{2,3}) e_3 + \varphi_{1,2} e_{1,2} + \varphi_{2,3} e_{2,3} - 2 w_2 e_{3,0} + 2\Lambda_1^2,\\ B_{1,1} &= u_3 K - \frac{\varphi_1 - t_{1,1}}{2} \alpha_0^\vee + \frac{\varphi_0 - t_{1,1}}{2} \alpha_1^\vee + \frac{w_2 \varphi_{1,2}}{2} \alpha_2^\vee + \frac{\varphi_{1,2}}{2} e_2 - w_2 e_3 + \Lambda_1,\end{aligned}$$

where

$$\begin{aligned}&\varphi_0 = 2w_1 + 2w_{2,3} + t_{1,1}, \quad \varphi_1 = -2w_0 - 2w_{2,3} + t_{1,1},\\ &\varphi_2 = (w_0 - 2w_1 + t_{1,1}) w_3 - 2w_{3,0}, \quad \varphi_3 = 2w_{1,2},\\ &\varphi_{1,2} = 2w_3, \quad \varphi_{2,3} = 2w_0 - 2w_1 + t_{1,1}.\end{aligned}$$

Note that $\kappa_0, \ldots, \kappa_4$ are constants. We also remark that

$$2w_2 \varphi_2 - \varphi_3 \varphi_{1,2} = 2(\kappa_2 - \kappa_3 - 3\rho_1), \quad \varphi_0 + \varphi_1 + \varphi_{2,3} = 3t_{1,1}.$$

Hence the system (6.3) can be expressed as a system of ordinary differential equations in terms of the variables $\varphi_0, \varphi_1, \varphi_2, \varphi_{1,2}, w_2$.

Let

$$q_1 = -\frac{w_2 \varphi_{1,2}}{\sqrt{6}}, \quad p_1 = -\frac{2\varphi_2}{\sqrt{6}\varphi_{1,2}}, \quad q_2 = \frac{\varphi_1}{\sqrt{6}}, \quad p_2 = -\frac{\varphi_0}{\sqrt{6}}, \quad t = -\frac{\sqrt{6} t_{1,1}}{2}.$$

We also set

$$\begin{aligned}&\alpha_1 = \frac{1}{3}(\kappa_2 - \kappa_3 - 3\rho_1), \quad \alpha_2 = \frac{1}{3}(\kappa_1 - 2\kappa_2 + \kappa_3),\\ &\alpha_3 = \frac{1}{3}(1 + \kappa_0 - 2\kappa_1 + \kappa_2), \quad \alpha_4 = \frac{1}{3}(1 - 2\kappa_0 + \kappa_1 + \kappa_3).\end{aligned}$$

Then we have

Theorem 6.2. *The system* (6.3) *with* (6.4) *gives the Painlevé system* $\mathcal{H}^{A_4^{(1)}}$. *Furthermore,* $\varphi_{1,2}$ *satisfies the completely integrable Pfaffian equation*

$$\frac{d}{dt} \log \varphi_{1,2} = p_1 + p_2 - \frac{2}{3} t.$$

6.3 For the partition $(4,1)$

The Heisenberg subalgebra $\mathfrak{s}_{(4,1)}$ of $\mathfrak{g}(A_4^{(1)})$ is defined by

$$\mathfrak{s}_{(4,1)} = \bigoplus_{k\in\mathbb{Z}\setminus 4\mathbb{Z}} \mathbb{C}\Lambda_1^k \oplus \bigoplus_{k\in\mathbb{Z}\setminus\{0\}} \mathbb{C}z^k H_1 \oplus \mathbb{C}K,$$

with

$$\Lambda_1 = e_0 + e_1 + e_4 + e_{2,3}, \quad H_1 = \alpha_1^\vee + 2\alpha_2^\vee - 2\alpha_3^\vee - \alpha_4^\vee.$$

The subalgebra $\mathfrak{s}_{(4,1)}$ admits the gradation of type $\mathbf{s} = (2,2,1,1,2)$ with the grade operator

$$\vartheta_{(4,1)} = 8\left(z\frac{d}{dz} + \mathrm{ad}\eta_{(4,1)}\right), \quad \eta_{(4,1)} = \frac{1}{8}(3\alpha_1^\vee + 4\alpha_2^\vee + 4\alpha_3^\vee + 3\alpha_4^\vee).$$

Note that

$$\mathfrak{g}_{\geq 0}(2,2,1,1,2) = \mathfrak{b}_+.$$

We now assume $t_{1,2} = 1$ and $t_{1,k} = 0$ $(k \geq 3)$. Then the similarity reduction (4.5) for $\mathfrak{s}_{(4,1)}$ is expressed as

$$\left[\vartheta_{(4,1)} - M, \partial_{1,1} - B_{1,1}\right] = 0, \tag{6.5}$$

with

$$\begin{aligned} M &= \vartheta_{(4,1)}(W)W^{-1} + W(\rho_1 H_1 + 2t_{1,1}\Lambda_1 + 4\Lambda_1^2)W^{-1}, \\ B_{1,1} &= \partial_{1,1}(W)W^{-1} + W\Lambda_1 W^{-1}. \end{aligned} \tag{6.6}$$

Let

$$W = \exp(\omega_{-1})\exp(\omega_{-2})\exp(\omega_{-3})\exp(\omega_{-4})\exp(\omega_{<-4}),$$

where

$$\begin{aligned} \omega_{-1} &= -w_2 f_2 - w_3 f_3, \\ \omega_{-2} &= -w_0 f_0 - w_1 f_1 - w_4 f_4 - w_{2,3} f_{2,3}, \\ \omega_{-3} &= -w_{1,2} f_{1,2} - w_{3,4} f_{3,4}, \\ \omega_{-4} &= -w_{0,1} f_{0,1} - w_{4,0} f_{4,0} - w_{1,2,3} f_{1,2,3} - w_{2,3,4} f_{2,3,4}, \end{aligned}$$

and $\omega_{<-4} \in \mathfrak{g}_{<-4}(2,2,1,1,2)$. Then the system (6.6) gives explicit formulas of $M, B_{1,1}$ as follows:

$$\begin{aligned} M = \;& \kappa_0\alpha_0^\vee + \kappa_1\alpha_1^\vee + \kappa_2\alpha_2^\vee + \kappa_3\alpha_3^\vee + \kappa_4\alpha_4^\vee + \varphi_0 e_0 + \varphi_1 e_1 \\ & + \varphi_2 e_2 + \varphi_3 e_3 + \varphi_4 e_4 + \varphi_{1,2} e_{1,2} + \varphi_{2,3} e_{2,3} + \varphi_{3,4} e_{3,4} + 4\Lambda_1^2, \\ B_{1,1} = \;& u_4 K + u_0\alpha_0^\vee + \frac{\varphi_0 - 2t_{1,1}}{4}\alpha_1^\vee + u_2\alpha_2^\vee + u_3\alpha_3^\vee + \frac{\varphi_{1,2}}{4}e_2 + \frac{\varphi_{3,4}}{4}e_3 + \Lambda_1, \end{aligned}$$

where

$$\begin{aligned}
&\varphi_0 = 4w_1 - 4w_4 + 2t_{1,1}, \quad \varphi_1 = -4w_0 + 2w_2w_3 - 4w_{2,3} + 2t_{1,1},\\
&\varphi_2 = -2(2w_1 - w_4 - t_{1,1})w_3 - 4w_{3,4}, \quad \varphi_3 = 2(w_1 - 2w_4 - t_{1,1})w_2 + 4w_{1,2},\\
&\varphi_{1,2} = 4w_3, \quad \varphi_{2,3} = -4w_1 + 4w_4 + 2t_{1,1}, \quad \varphi_{3,4} = -4w_2,
\end{aligned}$$

and

$$\begin{aligned}
64t_{1,1}u_0 &= (\varphi_0 - 4t_{1,1})(4\varphi_1 + \varphi_{1,2}\varphi_{3,4}) + 4\varphi_2\varphi_{3,4}\\
&\quad + 16t_{1,1}^2 + 16(\kappa_0 - \kappa_1 + \kappa_2 - \kappa_4 - 2\rho_1),\\
64t_{1,1}u_2 &= \varphi_0(4\varphi_1 + \varphi_{1,2}\varphi_{3,4}) + 4(\varphi_2 - t_{1,1}\varphi_{1,2})\varphi_{3,4}\\
&\quad - 16t_{1,1}^2 + 16(\kappa_0 - \kappa_1 + \kappa_2 - \kappa_4 - 2\rho_1),\\
64t_{1,1}u_3 &= \varphi_0(4\varphi_1 + \varphi_{1,2}\varphi_{3,4}) + 4\varphi_2\varphi_{3,4}\\
&\quad - 16t_{1,1}^2 + 16(\kappa_0 - \kappa_1 + \kappa_2 - \kappa_4 - 2\rho_1).
\end{aligned}$$

Note that $\kappa_0, \dots, \kappa_4$ are constants. We also remark that

$$\begin{aligned}
&(\varphi_0 - 4t_{1,1})\varphi_{1,2}\varphi_{3,4} + 4\varphi_3\varphi_{1,2} + 4\varphi_2\varphi_{3,4} = 16(-\kappa_2 + \kappa_3 + 4\rho_1),\\
&4\varphi_1 + 4\varphi_4 + \varphi_{1,2}\varphi_{3,4} = 16t_{1,1}, \quad \varphi_0 + \varphi_{2,3} = 4t_{1,1}.
\end{aligned}$$

Hence the system (6.5) can be described as a system of ordinary differential equations in terms of the variables $\varphi_0, \varphi_1, \varphi_2, \varphi_{1,2}, \varphi_{3,4}$.

Let

$$\begin{aligned}
&q_1 = \frac{\varphi_0}{4t_{1,1}}, \quad p_1 = \frac{t_{1,1}\varphi_1}{8},\\
&q_2 = \frac{\varphi_0}{4t_{1,1}} + \frac{\varphi_2}{t_{1,1}\varphi_{1,2}}, \quad p_2 = \frac{t_{1,1}\varphi_{1,2}\varphi_{3,4}}{32}, \quad t = -\frac{t_{1,1}^2}{2}.
\end{aligned}$$

We also set

$$\begin{aligned}
&\alpha_1 = \frac{1}{8}(2 - 2\kappa_0 + \kappa_1 + \kappa_4), \quad \alpha_2 = \frac{1}{8}(2 + \kappa_0 - 2\kappa_1 + \kappa_2),\\
&\alpha_3 = \frac{1}{8}(1 + \kappa_1 - 2\kappa_2 + \kappa_3), \quad \alpha_4 = \frac{1}{8}(\kappa_2 - \kappa_3 - 4\rho_1),\\
&\alpha_5 = \frac{1}{8}(1 - \kappa_3 + \kappa_4 + 4\rho_1).
\end{aligned}$$

Then we have

Theorem 6.3. *The system* (6.5) *with* (6.6) *gives the Painlevé system* $\mathcal{H}^{A_5^{(1)}}$. *Furthermore,* $\varphi_{1,2}$ *satisfies the completely integrable Pfaffian equation*

$$t\frac{d}{dt}\log\varphi_{1,2} = -q_1p_1 - q_2p_2 + tq_2 - \frac{3}{4}t - \frac{1 + 2\alpha_1 + 2\alpha_3 + 2\alpha_5}{4}.$$

A Lax pair

In the previous section, we have derived several Painlevé systems. Each of them can be regarded as the compatibility condition of a Lax pair (see Remark 4.1)

$$\frac{d\Psi}{dt} = B\Psi, \quad \vartheta_{\mathbf{n}}(\Psi) = M\Psi.$$

In this section, we give an explicit description of M and B by means of a bases of $\mathfrak{sl}_{n+1}[z, z^{-1}]$.

A.1 For the partition $(2, 2)$

The matrix M is described as follows:

$$M = \begin{bmatrix} \varepsilon_1 & -\frac{2(qp+\alpha_1+\alpha_2)}{w_1} & \sqrt{t} & 0 \\ 0 & \varepsilon_2 & \frac{w_1(q-t)}{\sqrt{t}} & 1 \\ \sqrt{t}z & 0 & \varepsilon_3 & \frac{2\sqrt{t}p}{w_1} \\ w_1(1-q)z & z & 0 & \varepsilon_4 \end{bmatrix},$$

where $\varepsilon_1, \dots, \varepsilon_4$ are linear conbinations of $\alpha_0, \dots, \alpha_3$. The matrix B is expressed as follows:

$$B = \frac{1}{2\sqrt{t}} \begin{bmatrix} u_1 - u_0 & x_1 & 1 & 0 \\ 0 & u_2 - u_1 & x_2 & 0 \\ z & 0 & u_3 - u_2 & x_3 \\ x_0 z & 0 & 0 & u_0 - u_3 \end{bmatrix}.$$

Each component of B is rational in q, p, w_1; see Section 6.1. The compatibility condition of this Lax pair gives the sixth Painlevé equation.

Remark A.1. *It is known that P_{VI} arises from the Lax pairs of two types, 2×2 matrix system* [IKSY] *and 8×8 matrix system* [NY3]*. The result of this section means that we derive a new Lax pair for P_{VI}.*

A.2 For the partition $(3, 1)$

The matrix M is described as follows:

$$M = \begin{bmatrix} \varepsilon_1 & \sqrt{6}(q_2 - q_1) & \varphi_{1,2} & 2 \\ 2z & \varepsilon_2 & -\frac{\sqrt{6}\varphi_{1,2}p_1}{2} & \sqrt{6}(p_2 - q_2 - t) \\ \frac{2\sqrt{6}q_1}{\varphi_{1,2}}z & 0 & \varepsilon_3 & \frac{6\{q_1(p_1+p_2-q_2-t)-\alpha_1\}}{\varphi_{1,2}} \\ -\sqrt{6}p_2 z & 2z & 0 & \varepsilon_4 \end{bmatrix},$$

where $\varepsilon_1, \dots, \varepsilon_4$ are linear conbinations of $\alpha_0, \dots, \alpha_3$. The matrix B is expressed as follows:

$$B = \frac{-2}{\sqrt{6}} \begin{bmatrix} u_1 - u_0 & 1 & 0 & 0 \\ 0 & u_2 - u_1 & x_2 & 1 \\ 0 & 0 & u_3 - u_2 & x_3 \\ z & 0 & 0 & u_0 - u_3 \end{bmatrix}.$$

Each component of B is rational in $q_1, p_1, q_2, p_2, \varphi_{1,2}$; see Section 6.2. The compatibility condition of this Lax pair gives the Painlevé system $\mathcal{H}^{A_4^{(1)}}$.

Note that the system $\mathcal{H}^{A_4^{(1)}}$ also arise from the Lax pair by means of 5×5 matricies [NY1].

A.3 For the partition $(4, 1)$

The matrix M is described as follows:

$$M = \begin{bmatrix} \varepsilon_1 & \frac{8p_1}{\sqrt{-2t}} & \varphi_{1,2} & 4 & 0 \\ 0 & \varepsilon_2 & \sqrt{-2t}\varphi_{1,2}(q_2 - q_1) & 4\sqrt{-2t}(1 - q_1) & 4 \\ 0 & 0 & \varepsilon_3 & \frac{32\{(1-q_2)p_2-\alpha_4\}}{\varphi_{1,2}} & \frac{32p_2}{\sqrt{-2t}\varphi_{1,2}} \\ 4z & 0 & 0 & \varepsilon_4 & -\frac{8(p_1+p_2+t)}{\sqrt{-2t}} \\ 4\sqrt{-2t}q_1 z & 4z & 0 & 0 & \varepsilon_5 \end{bmatrix},$$

where $\varepsilon_1, \dots, \varepsilon_5$ are linear conbinations of $\alpha_0, \dots, \alpha_4$. The matrix B is expressed as follows:

$$B = \frac{1}{\sqrt{-2t}} \begin{bmatrix} u_1 - u_0 & 1 & 0 & 0 & 0 \\ 0 & u_2 - u_1 & x_2 & 1 & 0 \\ 0 & 0 & u_3 - u_2 & x_3 & 0 \\ 0 & 0 & 0 & u_4 - u_3 & 1 \\ z & 0 & 0 & 0 & u_0 - u_4 \end{bmatrix}.$$

Each component of B is rational in $q_1, p_1, q_2, p_2, \varphi_{1,2}$; see Section 6.3. The compatibility condition of this Lax pair gives the Painlevé system $\mathcal{H}^{A_5^{(1)}}$.

Note that the system $\mathcal{H}^{A_5^{(1)}}$ also arise from the Lax pair by means of 6×6 matricies [NY1].

A.4 For the partition $(2,2,1)$

The matrix M is described as follows:

$$M=\begin{bmatrix} 0 & -\frac{4\sqrt{t}\varphi_{3,4}p_2}{\varphi_3} & \frac{8\sqrt{t}(q_1p_1+q_2p_2+\eta)}{\varphi_3} & 2\sqrt{t} & 0 \\ 0 & 0 & \varphi_2 & \frac{2\varphi_3(tq_2-q_1)}{t\varphi_{3,4}} & 2 \\ 0 & 0 & 0 & \frac{\varphi_3(t-q_1)}{s} & \varphi_{3,4} \\ 2\sqrt{t}z & 0 & 0 & 0 & -\frac{4t\varphi_{3,4}p_1}{\varphi_3} \\ \frac{2\varphi_3(q_1-q_2)}{\sqrt{t}\varphi_{3,4}}z & 2z & 0 & 0 & 0 \end{bmatrix},$$

where $\varepsilon_1,\dots,\varepsilon_5$ are linear conbinations of $\alpha_0,\dots,\alpha_4$ and

$$\varphi_2=\frac{8\{(q_2-1)(q_1p_1+q_2p_2+\eta)+\alpha_3\}}{\varphi_{3,4}}.$$

The matrix B is expressed as follows:

$$B=\frac{1}{2\sqrt{t}}\begin{bmatrix} u_1-u_0 & x_1 & x_{1,2} & 1 & 0 \\ 0 & u_2-u_1 & x_2 & x_{2,3} & 0 \\ 0 & 0 & u_3-u_2 & x_3 & 0 \\ z & 0 & 0 & u_4-u_3 & x_4 \\ x_0z & 0 & 0 & 0 & u_0-u_4 \end{bmatrix}.$$

Each component of B is rational in $q_1,p_1,q_2,p_2,\varphi_3,\varphi_{3,4}$; see Section 5.2. The compatibility condition of this Lax pair gives the system (1.1) with (1.2).

A.5 For the partition $(3,3)$

The matrix M is described as follows:

$$M=\begin{bmatrix} \varepsilon_1 & \frac{3t^{2/3}p_1}{w_3} & \frac{1}{t^{1/3}} & 0 & 0 & 0 \\ 0 & \varepsilon_2 & \frac{w_3(t-q_1)}{t} & 1 & 0 & 0 \\ 0 & 0 & \varepsilon_3 & -\frac{3(q_1p_1+q_2p_2+\eta)}{w_3} & \frac{1}{t^{1/3}} & 0 \\ 0 & 0 & 0 & \varepsilon_4 & \frac{w_3(q_2-1)}{t^{1/3}} & 1 \\ \frac{1}{t^{1/3}}z & 0 & 0 & 0 & \varepsilon_5 & \frac{3t^{1/3}p_2}{w_3} \\ \frac{w_3(q_1-q_2)}{t^{2/3}}z & z & 0 & 0 & 0 & \varepsilon_6 \end{bmatrix},$$

where $\varepsilon_1,\dots,\varepsilon_6$ are linear conbinations of $\alpha_0,\dots,\alpha_5$. The matrix B is expressed as follows:

$$B=\frac{-1}{3t^{4/3}}\begin{bmatrix} u_1-u_0 & x_1 & 1 & 0 & 0 & 0 \\ 0 & u_2-u_1 & x_2 & 0 & 0 & 0 \\ 0 & 0 & u_3-u_2 & x_3 & 1 & 0 \\ 0 & 0 & 0 & u_4-u_3 & x_4 & 0 \\ z & 0 & 0 & 0 & u_5-u_4 & x_5 \\ x_0z & 0 & 0 & 0 & 0 & u_0-u_5 \end{bmatrix}.$$

Each component of B is rational in q_1, p_1, q_2, p_2, w_3; see Section 5.1. The compatibility condition of this Lax pair gives the system (1.1) with (1.2).

B Affine Weyl group symmetry

The system (1.1) with (1.2) admits affine Weyl group symmetry of type $A_5^{(1)}$. In this section, we describe its action on the dependent variables and parameters.

Let r_i $(i = 0, \ldots, 5)$ be birational canonical transformations defined by

$$\begin{aligned} &\alpha_0 \to -\alpha_0, \quad \alpha_1 \to \alpha_0 + \alpha_1, \quad \alpha_5 \to \alpha_0 + \alpha_5, \\ &p_1 \to p_1 - \frac{\alpha_0}{q_1 - q_2}, \quad p_2 \to p_2 - \frac{\alpha_0}{q_2 - q_1}, \end{aligned}$$

for $i = 0$;

$$\alpha_0 \to \alpha_0 + \alpha_1, \quad \alpha_1 \to -\alpha_1, \quad \alpha_2 \to \alpha_1 + \alpha_2, \quad q_1 \to q_1 + \frac{\alpha_1}{p_1},$$

for $i = 1$;

$$\alpha_1 \to \alpha_1 + \alpha_2, \quad \alpha_2 \to -\alpha_2, \quad \alpha_3 \to \alpha_2 + \alpha_3, \quad p_1 \to p_1 - \frac{\alpha_2}{q_1 - t},$$

for $i = 2$;

$$\begin{aligned} &\alpha_2 \to \alpha_2 + \alpha_3, \quad \alpha_3 \to -\alpha_3, \quad \alpha_4 \to \alpha_3 + \alpha_4, \\ &q_1 \to q_1 + \frac{\alpha_3 q_1}{q_1 p_1 + q_2 p_2 - \alpha_3 + \eta}, \quad p_1 \to p_1 - \frac{\alpha_3 p_1}{q_1 p_1 + q_2 p_2 + \eta}, \\ &q_2 \to q_2 + \frac{\alpha_3 q_2}{q_1 p_1 + q_2 p_2 - \alpha_3 + \eta}, \quad p_2 \to p_2 - \frac{\alpha_3 p_2}{q_1 p_1 + q_2 p_2 + \eta}, \end{aligned}$$

for $i = 3$;

$$\alpha_3 \to \alpha_3 + \alpha_4, \quad \alpha_4 \to -\alpha_4, \quad \alpha_5 \to \alpha_4 + \alpha_5, \quad p_2 \to p_2 - \frac{\alpha_4}{q_2 - 1},$$

for $i = 4$;

$$\alpha_0 \to \alpha_0 + \alpha_5, \quad \alpha_4 \to \alpha_4 + \alpha_5, \quad \alpha_5 \to -\alpha_5, \quad q_2 \to q_2 + \frac{\alpha_5}{p_2},$$

for $i = 5$. Then the system (1.1) with (1.2) is invariant under the action of them. Furthermore, a group of symmetries $\langle r_0, \ldots, r_5 \rangle$ is isomorphic to the affine Weyl group of type $A_5^{(1)}$.

The group of symmetries defined above arises from the gauge transformations

$$r_i(\Psi) = \exp\left(\frac{\alpha_i}{\varphi_i} f_i\right)\Psi \quad (i = 0, \dots, 5),$$

where

$$\varphi_0 = \frac{w_3(q_2 - q_1)}{3t^{2/3}}, \quad \varphi_1 = -\frac{t^{2/3} p_1}{w_3}, \quad \varphi_2 = \frac{w_3\,(q_1 - t)}{3t},$$
$$\varphi_3 = \frac{q_1 p_1 + q_2 p_2 + \eta}{w_3}, \quad \varphi_4 = \frac{w_3(1 - q_2)}{3t^{1/3}}, \quad \varphi_5 = -\frac{t^{1/3} p_2}{w_3},$$

for the Lax pair of Appendix A.5. Note that those transformations are derived from the following ones [NY2]:

$$r_i(G) = G \exp(-e_i)\exp(f_i)\exp(-e_i) \quad (i = 0, \dots, 5),$$

where G is an $N_- B_+$-valued function given in Section 4.

Acknowledgement

The authers are grateful to Professors Kenji Kajiwara, Masatoshi Noumi, Masahiko Saito and Yasuhiko Yamada for valuable discussions and advices.

References

[AS] M. J. Ablowitz and H. Segur, Exact linearization of a Painlevé transcendent, Phys. Rev. Lett. **38** (1977) 1103-1106.

[C] R. Carter, Conjugacy classes in the Weyl group, Compositio Math. **25** (1972), 1-59.

[DF] F. Delduc and L. Fehér, Regular conjugacy classes in the Weyl group and integral hierarchies, J. Phys. A: Math. Gen. **28** (1995) 5843-5882.

[DS] V. G. Drinfel'd and V. V. Sokolov, Lie algebras and equations of Korteweg-de Vries type, J. Sov. Math. **30** (1985) 1975-2036.

[FS] K. Fuji and T. Suzuki, Higher order Painlevé system of type $D^{(1)}_{2n+2}$ arising from integrable hierarchy, Int. Math. Res. Not. **1** (2008) 1-21.

[GHM] M. F. de Groot, T. J. Hollowood and J. L. Miramontes, Generalized Drinfeld-Sokolov hierarchies, Comm. Math. Phys. **145** (1992) 57-84.

[IKSY] K. Iwasaki, H. Kimura, S. Shimomura and M. Yoshida, From Gauss to Painlevé — A Modern Theory of Special Functions, Aspects of Mathematics **E16** (Vieweg, 1991).

[Kac] V. G. Kac, Infinite dimensional Lie algebras, Cambridge University Press (1990).

[KIK] T. Kikuchi, T. Ikeda and S. Kakei, Similarity reduction of the modified Yajima-Oikawa equation, J. Phys. A: Math. Gen., **36** (2003) 11465-11480.

[KK1] S. Kakei and T. Kikuchi, Affine Lie group approach to a derivative nonlinear Schrödinger equation and its similarity reduction, Int. Math. Res. Not. **78** (2004) 4181-4209;

[KK2] S. Kakei and T. Kikuchi, The sixth Painlevé equation as similarity reduction of $\widehat{\mathfrak{gl}}_3$ hierarchy, Lett. Math. Phys. **79** (2007) 221-234.

[KL] F. ten Kroode and J. van de Leur, Bosonic and fermionic realizations of the affine algebra $\widehat{\mathfrak{gl}}_n$, Comm. Math. Phys. **137** (1991) 67-107.

[KP] V. G. Kac and D. Peterson, 112 constructions of the basic representation of the roop group of E_8, in Symposium on Anomalies, Geometry ans Topology, ed. W. A. Baedeen and A. R. White, (World Scientific, 1985) 276-298.

[NY1] M. Noumi and Y. Yamada, Higher order Painlevé equations of type $A_l^{(1)}$, Funkcial. Ekvac. **41** (1998), 483-503.

[NY2] M. Noumi and Y. Yamada, Birational Weyl group action arising from a nilpotent Poisson algebra, in Physics and Combinatorics 1999, Proceedings of the Nagoya 1999 International Workshop, ed. A.N.Kirillov, A.Tsuchiya and H.Umemura, (World Scientific, 2001) 287-319.

[NY3] M. Noumi and Y. Yamada, A new Lax pair for the sixth Painlevé equation associated with $\widehat{\mathfrak{so}}(8)$, in Microlocal Analysis and Complex Fourier Analysis, ed. T.Kawai and K.Fujita, (World Scientific, 2002) 238-252.

[S] Y. Sasano; Higher order Painlevé equations of type $D_l^{(1)}$, RIMS Koukyuroku **1473** (2006) 143-163.